

National and Kapodistrian University of Athens

Faculty of Pharmacy
Department of Pharmacognosy
& Natural Products Chemistry
PanepistimiopolisZografou
15 771 Athens
Tel:+30 210 72 74052
magiatis@pharm.uoa.gr



Athens, 29/12/2016 N°:485/2016

CERTIFICATE OF ANALYSIS

WILLIAM TO

Owner: Sakellaropoulos Organic Farming

Origin: Sparta Lakonia, Greece **Harvest season**: November 2016

Variety: Koroneiki

Sample: Armonia organic EVOO

Physical properties: significant pungent and moderate bitter character

Chemical analysis

Oleocanthal: 223 mg/Kg Oleacein: 90 mg/Kg

Oleuropein aglycon (monoaldehyde form): 50 mg/Kg Oleuropein aglycon (dialdehyde forms)*: 32 mg/Kg Ligstroside aglycon (monoaldehyde form): 48 mg/Kg Ligstroside aglycon (dialdehyde forms)**: 96 mg/Kg

Total hydroxytyrosol derivatives: 172 mg/Kg Total derivatives of tyrosol: 367 mg/Kg Oleocanthal+Oleacein (Index D1): 313 mg/Kg

Total of analyzed compounds (index D3): 539 mg/Kg

Comments

The daily consumption of 20 g of the analyzed olive oil sample provides 10.8 mg of hydroxytyrosol, tyrosol or their derivatives (>5 mg) and consequently the oil belongs to the category of oils that protect the blood lipids from oxidative stress according to the Regulation 432/2012 of the European Union.

The level of oleocanthal is higher than the average value (135 mg/Kg) of the samples included in the international study performed at the University of California, Davis.

It should be noted that oleocanthal and oleacein present important biological activity and they have been related with anti-inflammatory, antioxidant, cardioprotective and neuroprotective activity.

The chemical analysis was performed according to the method published in J. Agric. Food Chem., 2012, 60 (47), pp 11696–11703, J. Agric. FoodChem., 2014, 62(3), 600–607 and OLIVAE, 2015, 122, 22-33.

*Oleomissional+Oleuropeindial**Ligstrodial+Oleokoronal

Prokopios Magiatis

PROKOFIOS MAGIATIS
ASSOCIATE PROFESSOR
UNIVERSAL OF ATHEMS
FACULTY OF PHARMACY
DEPARTMENT OF PHARMACOGNOSY
AND NATURAL PRODUCTS CHEMISTRY