

## National and Kapodistrian University of Athens

Faculty of Pharmacy
Department of Pharmacognosy & Natural Products Chemistry
Panepistimiopolis Zografou
15771, Athens
Tel: +30 210 72 74052



Athens, 27/12/2017 Cert.Num: 1718-C00428

## **CERTIFICATE OF ANALYSIS**

Brand Name: ARMONIA EVOO Analysis Date: 27/12/2017

Owner: SAKELLAROPOULOS ORGANIC FARMING

**Origin:** SPARTI LAKONIA - GREECE

## **Chemical Analysis**

magiatis@pharm.uoa.gr

Oleocanthal	231	mg/Kg
Oleacein	152	mg/Kg
Oleocanthal + Oleacein (index D1)	383	mg/Kg
Ligstroside aglycon (monoaldehyde form)	58	mg/Kg
Oleuropein aglycon (monoaldehyde form)	38	mg/Kg
Ligstroside aglycon (dialdehyde form)	< 5	mg/Kg
Oleuropein aglycon (dialdehyde form)	< 5	mg/Kg
Total tyrosol derivatives	289	mg/Kg
Total hydroxytyrosol derivatives	190	mg/Kg
Total phenols analyzed	479	mg/Kg

## **Comments:**

The levels of oleocanthal and oleacein are higher than the average values (135 and 105 mg/Kg respectively) of the sample included in the international study performed at the University of California, Davis.

The daily consumption of 20 g of the analyzed olive oil provides 9.6 mg of hydroxytyrosol, tyrosol or their derivatives (>5 mg) and consequently the oil belongs to the category of oils that protect the blood lipids from oxidative stress according to the Regulation 432/2012 of the European Union.

It should be noted that oleocanthal and oleacein present important biological activity and they have been related with anti-inflammatory, antioxidant, cardioprotective and neuroprotective activity.

The chemical analysis was performed according to the method published in J.Agric. Food Chem., 2012, 60 (47), pp 11696-11703, J.Agric. Food Chem., 2014 62 (3), 600-607 and OLIVAE, 2015, 122, 22-33.

\*Oleomissional+Oleuropeindial \*\*Ligstrodial+Oleokoronal

Magiatis Prokopios

PROKOPIOS MAGIATIS
ASSOCIATE PROFESSOR

UNIVERSITY OF ATHEN

DEPARTMENT OF HARMACOGNOSY
AND NATURAL PROPERTY OF HARMACOGNOSY
AND NATURAL PROPERTY OF HARMACOGNOSY