

National and Kapodistrian University of Athens

Faculty of Pharmacy
Department of Pharmacognosy & Natural Products Chemistry
Panepistimiopolis Zografou
15771, Athens
Tel: +30 210 72 74052



Athens, 27/12/2017 Cert.Num: 1718-C00426

Analysis Date: 27/12/2017

CERTIFICATE OF ANALYSIS

Brand Name: FYLLIKON Early Harvest EVOO

SAKELLAROPOULOS ORGANIC FARMING

Origin: SPARTI LAKONIA - GREECE

Chemical Analysis

magiatis@pharm.uoa.gr

Owner:

Oleocanthal	309	mg/Kg
Oleacein	230	mg/Kg
Oleocanthal + Oleacein (index D1)	540	mg/Kg
Ligstroside aglycon (monoaldehyde form)	58	mg/Kg
Oleuropein aglycon (monoaldehyde form)	69	mg/Kg
Ligstroside aglycon (dialdehyde form)	77	mg/Kg
Oleuropein aglycon (dialdehyde form)	52	mg/Kg
Total tyrosol derivatives	444	mg/Kg
Total hydroxytyrosol derivatives	351	mg/Kg
Total phenols analyzed	795	mg/Kg

Comments:

The levels of oleocanthal and oleacein are higher than the average values (135 and 105 mg/Kg respectively) of the sample included in the international study performed at the University of California, Davis.

The daily consumption of 20 g of the analyzed olive oil provides 15.9 mg of hydroxytyrosol, tyrosol or their derivatives (>5 mg) and consequently the oil belongs to the category of oils that protect the blood lipids from oxidative stress according to the Regulation 432/2012 of the European Union.

It should be noted that oleocanthal and oleacein present important biological activity and they have been related with anti-inflammatory, antioxidant, cardioprotective and neuroprotective activity.

The chemical analysis was performed according to the method published in J.Agric. Food Chem., 2012, 60 (47), pp 11696-11703, J.Agric. Food Chem., 2014 62 (3), 600-607 and OLIVAE, 2015, 122, 22-33.

*Oleomissional+Oleuropeindial **Ligstrodial+Oleokoronal

Magiatis Prokopios

PROKOPIOS MAGIATIS

ASSOCIATE PROFESSOR

FACULT DY PHARMACY
DEPARTMENT OF PHARMACOGNOSY
AND NATURAL PHONE CHEMISTRY