

National and Kapodistrian University of Athens

Faculty of Pharmacy Department of Pharmacognosy & Natural Products Chemistry Panepistimiopolis Zografou 15 771 Athens Tel:+30 210 72 74052 magiatis@pharm.uoa.gr



Athens, 9/11/2018 N°: 733/2018

CERTIFICATE OF ANALYSIS

Owner: SAKELLAROPOULOS OLIVE GROVES

Geographic origin: Lakonia, Greece

Chemical analysis

Name Tyrosol μg/g Hydroxytyrosol μg/g*

LEMONENIES 520 1180

Comments

The levels of tyrosol are higher than the average values of commercial olives samples (134 μ g/g)that were included in the study performed at the University of Athens and published in J. Agric. Food Chem. 2010, 58, 46–50. Oleuropein was not detected (<5 μ g/g).

It should be noted that hydroxytyrosol and tyrosol present important biological activity and they have been related with antioxidant and cardioprotective activity.

Daily consumption of 3 gr of the olives of this sample offers >5 mg of hydroxytyrosol and tyrosol and corresponds to the consumption of 20 gr of olive oil belonging to the oil category that protect the blood lipids from oxidative stress, according to the EU regulation 432/2012.

Prokopios Magiatis

*The values are expressed per wet weight of olive flesh

PROKOPIOS MAGIATIS
ASSOCIATE PROFESSOR
UNIVERSAL OF ATHEMS
FACULT OF PHARMACY
DEPARTMENT OF PHARMACOGNOSY
AND NATURAL PRODUCTS CHEMISTRY